

INTRA-OFFICE MEMORANDUM

25X1X6
 TO: [REDACTED]
 FROM: [REDACTED]
 SUBJECT: New Products - April, 1953

DATE: May 4, 1953

DomesticBicillin with Sulfase Tablets

Wyeth has now introduced Bicillin with Sulfa in a tablet form. It is supplied in bottles of 36 tablets. For a full description, see BiCillin Sulfas, New Products, November, 1952.

Bacitracin Ointment

Indications: In the prevention of infection following minor cuts and abrasions.
 Administration: Apply 1 or 2 times daily.
 Form Supplied: In tubes of $\frac{1}{2}$ oz.

25X1X6

Bacitracin Parenteral

Description: Each vial contains: 50,000 units
 Bacitracin
 Indications: Against a broad range of gram-positive organisms and penicillin-resistant strains.
 Administration: For hospitalized patients only, amount as determined by physician.
 Form Supplied: In vials as a dry powder.
 Source: 25X1X6 [REDACTED]

Diatracin

Description: Each gram contains: 500 units
 Bacitracin 0.05 Gm.
 Sulfadiazine
 Form Supplied: Topical Ointment - tubes of $\frac{1}{2}$ oz.
 Ophthalmic Ointment $\frac{1}{8}$ oz. tubes
 Source: 25X1X6 [REDACTED]

Dicrysticin

Description: The formula for this product has been changed from:
 Procaine Penicillin G 300,000 units
 Buffered Crystalline Penicillin
 G Potassium 100,000 units
 Dihydrostreptomycin Sulfate 0.5 Gm
 To:
 Procaine Penicillin G 300,000 units
 Potassium Penicillin G 100,000 units
 Streptomycin Sulfate 0.25 Gm.
 Dihydrostreptomycin Sulfate 0.25 Gm.

-2-

Dicrysticin Fortis

This dosage form contains the same amounts of penicillin plus 0.5 Gm. each of Streptomycin and Dihydrostreptomycin Sulfate.

Distrycillin - aqueous suspension

25X1X6	Procaine Penicillin G	400,000 units
	Streptomycin Sulfate	0.25 Gm.
	Dihydrostreptomycin Sulfate	0.25 Gm.

Dihydro Dispolator
Description:

Each dispolator contains 50 mg. micro-pulverized dihydrostreptomycin sulfate in disposable plastic inhalor.

Indications:

Recommended for local treatment of infections of the respiratory tract.

Administration:

2 Dispolators three times a day

Form Supplied:

2 vials, each containing 3 dispolators

Source:

Duo-Strep 25X1X6
Description:

A white powder containing the equivalent of 1 Gm. free base of streptomycin sulfate and dihydrostreptomycin sulfate in equal parts.

Indications:

It is indicated in the treatment of pulmonary, meningeal, miliary, genito-urinary, skeletal, serosal tuberculosis and tuberculous adenitis. It can be used where protracted treatment is necessary as toxicity is greatly reduced.

Administration:

As in the case of Streptomycin or Dihydrostreptomycin, it is best used in conjunction with another anti-TB drug such as PAS or Isoniazid.

Form Supplied:

25X1X6

Vials containing the equivalent of 1 gm. free base of streptomycin and dihydrostreptomycin sulfate.

Source:

Neomycin Ointment, Neomycin Ophthalmic Ointment

Description:

Both ointments contain:

Neomycin Sulfate

5 mg. per Gm.

Indications:

Impetigo, wound infections, sycosis barbae, conjunctivitis, burns, styes, boils, etc.

Administration:

Applied 3 or 4 times daily.

Form Supplied:

Ointment - Tubes $\frac{1}{2}$ oz., 1 oz., 4 oz.

Ophthalmic Ointment - $\frac{1}{8}$ oz. tubes

Source:

Eli Lilly & Co., Indianapolis, Indiana

Neomycin and Kaolin
Description:

This product is an aqueous preparation of neomycin sulfate, with kaolin and pectin suspended with methycellulose.

Neomycin and Kaolin cont'd.

Indications: Designed for the symptomatic treatment of diarrhea and as a specific therapy in bacterial diarrhea caused by neomycin-susceptible organisms.

25X1X6

Source: [REDACTED]

Panbiotic

Description: Each dose contains:
Procaine Penicillin G, Cryst. 300,000 units
Potassium Penicillin G 300,000 units
DBED Penicillin 600,000 units
Buffering agent in dry form

Form Supplied: 1-dose vials containing 1.2 million units
5-dose vials containing 6.0 million units

Source: 25X1X6 [REDACTED] New York

Pansulfa with Penicillin - New Package Size

Description: This product combines penicillin with triple-sulfonamides.

Indications: For treatment of infections which respond to oral administration of penicillin and triple-sulfonamide combinations.

Administration: As determined by the physician

Form Supplied: Now available in bottles of 100 tablets
25X1X6 as well as in bottles of 24 and 30

Source: [REDACTED]

Pastillas Antibioticas or Pastillas PSS

Description: Each lozenge contains:
Calcium Penicillin 1,000 U.I.
Streptomycin 2,000 U. (2 mg.)
Sulfadiazine 100 mg.

Indications: Used in treating infections of the gums, mouth and throat caused by Gram-negative or Gram-positive organisms sensitive to these antibiotics. Used also in operations on the oral cavity.

Administration: Take one lozenge and allow it to dissolve slowly in the mouth to permit contact with the infected area for as long as possible.

Source: 25X1X6 [REDACTED] AMIO

Redutyl

Description: Each vial contains:
INH 25X1X6 1 Gm.
Dihydrostreptomycin .5 Gm.

Form Supplied: 1 dose vials

Source: [REDACTED]
[REDACTED] in January).

-1-

Foreign

Atoximicina or Atralmicina

Description: The formula is as follows:
 Procaine Penicillin G 300,000 units
 Potassium Penicillin 100,000 units
 Streptomycin (Sulfate) 0.25 g.
 Dihydrostreptomycin 0.25 g.
 Source: 25X1X6 Paolo Cocco - Atoximicina

Cloromisan Lauzier

Description: Capsule - each one containing:
 Chloramphenicol U.S.A. XIV 0.25 gm.
 Dihydrostreptomycin Sulfate (equivalent to streptomycin base) 0.05 gm.

Indications: The combination of these two drugs extends and strengthens the antibacterial spectrum of each.

Administration: Best results are obtained by administering 2 capsules a day - or approximately 50 mmg. per kilo of body weight. In case of a fever the second or third day, use 1 capsule every four to six hours for 6 to 8 days.

Form Supplied: Bottles of 12.

Description: Ointment
 Each tube contains:
 Chloramphenicol U.S.P. 1 gm.
 Dihydrostreptomycin Sulfate (as above) 0.05 g.
 Propylparaben 0.02 g.
 Lanolin 30 g.
 Vaseline 58.93 g.
 Alcohol

Indications: Used in treatment of skin lesions, dermatosis and wounds caused by Gram-negative bacteria or in cases where the cause of infection is not known.

Administration: Use sufficient ointment to cover the affected parts. Repeat as often as necessary.

Form Supplied: Tubes of 28 g.

Description: Suppositories
 Each suppository contains:
 Chloramphenicol 0.25 g.
 Dihydrostreptomycin Sulfate (as above) 0.05 g.
 Propylparaben 0.0004 g.
 Cocoa-butter

Indications: Same as the capsules but can be used where the patient cannot respond to oral treatment because of nausea or vomiting due to some gastro-intestinal infections or pertusis.

Administration: In case of whooping cough in children of six months- Use 1 suppository in divided doses for four days.

For children five years or older - Use 1 to 4

Approved For Release 1999/09/10 : CIA-RDP83-00423R000700410009-0

Form Supplied: Boxes of 6 suppositories.

Source:

-5-

Estrepto-talil Linner

Description: This contains:
 Streptomycin Sulfate 0.080 Gr.
 Sulfathiazole 0.300 Gr.
 Sulfadiazine 0.125 Gr.
 Sulfamerazine 0.125 Gr.
 Bacterial antigens (Enteric) 10 million

Indications: Enteric bacterial infections, both acute and chronic. Also pre- and post-operatively, particularly in abdominal surgery.

Administration: Initially 4 tablets and then continue with 2 tablets every 4 hours. In case of pronounced diarrhea the cream is recommended. Children 4 teaspoonfuls initially; continue with 2 teaspoonfuls every four hours.

Form Supplied: Cream - Bottles of 100 cc.
 Tablets - Tubes of 20

Source: 25X1X6 [REDACTED]

Estricilina

Description: Each vial contains:
 Dihydrostreptomycin Sulfate - equivalent to 0.5 g. of Streptomycin base.
 Procaine Penicillin G 300,000 units
 Penicillin G Crystalline 100,000 units

Form Supplied: 1 Bottle of 5 cm³ with 1 vial of sterile solution of Sodium Chlorate to 0.85g.%

Source: 25X1X6 [REDACTED]

Nebacetin

Description: Each gm or ml. contains:
 Neomycin 5 mg.
 Bacitracin 250 units

Form Supplied: Dry powder for aqueous solution -
 Bottles of 10 and 30 ml.
 Salve - ophthalmic ointment - Tubes of 5 g. and 20 g.

Source: 25X1X6 [REDACTED]

Ophtovit

Description: Penicillin 20,000 U.
 Streptomycin 0.10 gr.
 Vitamin A 10,000 U
 Sulfathiol 1 gr.

Source: 25X1X6 [REDACTED]

Indications: Infections of the eye

Form Supplied: Ointment in tubes of 5 gr.

Source: [REDACTED]

25X1X6

Neotrimeran - P

Neotrizine Savorets in Venezuela under the new trade name shown above.

-6-

Penichicle Senosiain

Description: Each piece contains:
Potassium Penicillin G, Crystalline 10,000 U.O.
Chicle, sugar, mint, c.b.p.

Indications: Used in treatment of mouth and throat infections due to germs sensitive to the action of Penicillin.
Acute and chronic catarrhal and amigdilites, Vincent's Angina, Stomatitis and Gingivitis.

Administration: Chew one piece every 2 to 3 hours, according to the severity of the infection.

Form Supplied: 25X1X6 Boxes of 5, 10 and 20 tablets.

Source: [REDACTED] P.F.

Penisul

Description: Simple
White tablets, each one containing:
Sodium Penicillin Cryst. 5,833 Un.
Sulfadiazine 0.11 gr.
Talc, sugar, licorice and magnesium stearate.

Fortified
Sodium Penicillin G, Cryst. 150,000 Un.
Sulfadiazine 0.15 gr.
Sulfametazine 0.15 gr.
Excipient 0.6 gr.

Indications: Infections of the mouth and pharynx

Administration: Dissolve slowly in the mouth - 1 tablet every 2 hours. Doctor may prescribe other dosages.

Form Supplied: 25X1X6 Bottles of 25 tablets

Source: [REDACTED] la.

Rafacombin

Description: Each vial contains:
Streptomycin 1 gr.
Penicillin 400,000 units

Indications: Tuberculosis and infections due to Streptomycin and Penicillin sensitive organisms.

Administration: Intramuscular

Form Supplied: In vials 25X1X6

Source: [REDACTED]

Streptonazid

Description: Each vial contains:
Dihydrostreptomycin 1 g.
Hydrazide 0.25 g.

Indications: Tuberculosis

Form Supplied: Boxes of 1 vial

Source: [REDACTED] 25X1X6

Streptotibina

Description: This is the first molecular combination of dihydrostreptomycin and an isonicotinylnalidixic acid derivative.

Form Supplied: 25X1X6 Boxes of 1 vial

Source: [REDACTED]

-7-

Suerbional

Description: This product contains penicillin (either 500,000 or 1,000,000 units) and streptomycin.
Administration: 1 injection every 24 hours. If a larger dose is needed a second injection can be given within 12 to 16 to 18 hours.

25X1X6

Source:

Synergistine

Description: Each vial contains:
Streptomycin .5 Gr.
Penicillin 400,000 units
Indications: Tuberculosis and infections due to Streptomycin and Penicillin sensitive organisms.

25X1X6

Administration: Intramuscular

Source:

Trisul-Pen

Description: White tablets, each one containing:
Potassium Penicillin G 50,000 U.I.
Sulfadiazine 0.166 gr.
Sulfamerazine 0.166 gr.
Sulfametazine 0.166 gr.
Excipient

Indications: To be used according to doctor's instructions.

Administration: Orally as indicated by doctor.

Form Supplied: Bottles of 20 tablets

Source: Laboratories Pharmacia S.A. Caracas,

25X1X6

Tirobacin

Description: Each lozenge contains:
Bacitracin 1 mgr.
Tyrothrycin 1 mgr.
Benzocaine 2.5 mgr.
Excipient

Indications: Mouth and throat infections

Administration: Orally - to be chewed slowly

Form Supplied: Packages of 20 lozenges

Source:

25X1X6

Thiocillin Topical Ointment, Ophthalmic Ointment

Description: Each contains:
Penicillin 3,000 units
Sulfathiazole 0.50
Vitamin A 5,000 U.
Vitamin D 3,000 U.

Indications: Skin infections, eye infections

Form Supplied: Tubes of 20 gr. - topical ointment

Tubes of 8 gr. - ophthalmic ointment

Source:

25X1X6